

Together to End Stroke[™]

STROKE OCCURS IN TODDLERS, CHILDREN, AND TEENS. STROKE IS AN EMERGENCY AT ANY AGE!

KNOW?



IF ONE OR MORE OF THESE SIGNS ARE PRESENT, DON'T DELAY!

CALL 911 OR THE EMERGENCY SYSTEM IN YOUR COUNTRY!

IF IT LOOKS LIKE A STROKE

ADDITIONAL WARNING SIGNS IN CHILDREN

- Severe sudden headache, especially with vomiting and sleepiness
- Sudden weakness or numbness on one side of the body (face, arm and/or leg)
- Sudden confusion, difficulty speaking or understanding others
- Sudden trouble seeing to one side or loss of vision
- Sudden difficulty walking, dizziness, loss of balance or coordination
- New-onset of seizures usually on one side of the body

THINK STROKE

Signs of stroke are often missed in children and teens because there is a lack of awareness that strokes can happen in this age group.



PREVIOUSLY IDENTIFIED RISK FACTOR



TOP 10 CAUSES OF DEATH in children



The use of tPA (a clot busting treatment) or clot retrieval devices are

NOT FDA APPROVED FOR USE IN CHILDREN in the U.S.

International Alliance

for Pediatric Stroke

MEDICAL CONDITIONS ASSOCIATED Sickle Cell Disease, Moyamoya Syndrome, Arterial Dissection, WITH STROKE INCLUDE: Autoimmune Disorders, Congenital Heart Disease, Blood Clott

Autoimmune Disorders, Congenital Heart Disease, Blood Clotting Disorders

Learn more at iapediatricstroke.org and strokeassociation.org

Sources:

iapediatricstroke.org/infographic.pdf iapediatricstroke.org/2014%20AHA.fact.pdf

heart.org/idc/groups/heart-public/@wcm/@sop/@smd/documents/downloadable/ucm_472920.pdf